

3M™ Tegaderm™ Absorbent Clear Acrylic Dressing

Tegaderm™ Absorbent Clear Acrylic Dressing



The Difference
is Clear



3M™ Tegaderm™ Absorbent Clear Acrylic Dressing – a unique transparent wound dressing with fluid handling capacity

- Transparent dressing allows for wound monitoring without changing the dressing
- Clear design takes the guesswork out of application over the wound
- Novel acrylic polymer pad technology designed to handle low to moderate wound drainage

3M

3M™ Tegaderm™ Absorbent Clear Acrylic Dressing advances 3M™ Tegaderm™ Transparent Film Dressing technology by adding a clear acrylic polymer pad designed to handle low to moderate drainage.



TOP LAYER

- Breathable, letting oxygen in and moisture vapor out, allowing the skin to function normally
- Tegaderm™ Transparent Film barrier is impervious to liquids, bacteria, and viruses,¹ providing an effective barrier to external contaminants

MIDDLE LAYER

- Patented clear acrylic polymer pad handles exudate through the process of diffusion
- Handles low- to moderate-exuding wounds

BOTTOM LAYER

- Bottom layer of film is perforated to allow wound exudate to pass through the film to the acrylic polymer pad
- Unique adhesive designed for long wear, even on moist skin

Clearly Different

- Dressing designed not to melt down or leave residue
- Non-bulky design to help prevent dressing roll or lift due to friction from linens, clothing or prosthetics
- Eliminates hydrocolloid odor
- Allows for one-handed placement

Clinically Effective

- Low-friction surface minimizes potential for friction and shear
- Low potential for periwound maceration
- Maintains a moist wound healing environment
- Rated significantly better than the leading hydrocolloid dressing²
 - Ability to monitor the wound before and after absorption
 - Conformable before and after absorption
- Superior fluid handling capacity compared to leading hydrocolloid brands*

Affordable

- Without lifting or changing the dressing, easily monitor the wound for:
 - Color of drainage
 - Condition of periwound skin
 - Condition of wound bed
- Less frequent dressing changes may save nursing time and supply costs
 - Can be worn until the dressing leaks, loses adhesion or according to facility protocol

Comfortable for Patients

- Less frequent dressing changes reduce the risk of skin trauma
- Allows gentle removal from skin
- Washable – can easily wipe off stool, urine or other contaminants
- Waterproof – allows patients to bathe or shower*

*Data on file at 3M.

Where to use 3M™ Tegaderm™ Absorbent Clear Acrylic Dressing



Over skin tears and superficial traumatic wounds



As a primary dressing over Stage II/Stage III low- to moderate-draining pressure ulcers



As a primary dressing over skin graft donor sites



Over pressure-related injuries



As a secondary dressing cover over wounds requiring an absorbent wound filler



As a primary dressing over surgical wounds

Skin Tears: State of the Science: Consensus statements for the prevention, predication, assessment and treatment of skin tears¹²


Tegaderm™ Absorbent Clear Acrylic Dressings were listed as a successful treatment for categories 1 to 3 skin tears. The dressings were left in place for 21 days, minimal pain was reported upon removal and complete wound closure was seen in all cases.¹²



Skin Tear Treatment

Skin Tears occur when the epidermis separates from the dermis. They are often painful and distressing in appearance. They represent a major problem affecting older adults and many compromised individuals. It is estimated that at least 1.5 million skin tears occur in institutionalized older adults each year,⁴ with individual facilities reporting prevalence rates between 14% and 24%.⁵

Health Economic Data for Skin Tear Treatment*



- A 3M™ Tegaderm™ Absorbent Clear Acrylic Dressing^{3,6}
- B Mölnlycke Mepilex® R Border Silicone Foam Dressing^{6,7}
- C Smith & Nephew Flexigel™ Hydrogel Sheet Dressing^{7,8,9}
- D Systagenix ADAPTIC® Nonadhering Dressing^{8,9,10}
- E Kendall Telfa® "Ouchless" Non-Adherent Dressing^{5,7}

Includes:
 Standard supplies (3 pair gloves, wound cleanser, biohazard bag)
 Labor – 15 minutes¹¹
 Frequency of dressing change based on references
 *Data on file at 3M.



Visit YouTube to learn how 2 generations of nurses saw the difference Tegaderm™ Absorbent Clear Acrylic Dressing made in their lives







<http://YouTu.be/C99QiLxzdEw>



3M™ Tegaderm™ Absorbent Clear Acrylic Dressing

Ordering Information:

Dressing should overlap the wound margins by at least 3/8" (1 cm).

	Catalog Number	NDC/NHRC Number	Dressing Pad Size	Overall Dressing Size	Dressings/Box	Boxes/Case	HCPCS Code
	90800 Oval	8333-9080-00	1½ in. x 2¼ in. 3,8 x 5,7 cm	3 in. x 3¾ in. 7,6 x 9,5 cm	5	6	A6203
	90801 Oval	8333-9080-10	2¾ in. x 3 in. 6,0 x 7,6 cm	4¾ in. x 5 in. 11,1 x 12,7 cm	5	6	A6203
	90802 Square	8333-9080-20	3¾ in. x 4 in. 9,8 x 10,1 cm	5¾ in. x 6 in. 14,9 x 15,2 cm	5	6	A6203
	90803 Oval	8333-9080-30	3¾ in. x 4¼ in. 8,5 x 10,7 cm	5¾ in. x 6¼ in. 14,2 x 15,8 cm	5	6	A6203
	90805 Square	8333-9080-50	5¾ in. x 6 in. 14,9 x 15,2 cm	7¾ in. x 8 in. 20,0 x 20,3 cm	5	4	A6204
	90807 Sacral	8333-9080-70	4½ in. x 5 5/8 in. 11,7 x 14,3 cm	6 5/8 in. x 7 ½ in. 16,8 x 19 cm	5	4	A6204

Disclaimer:

The reimbursement information provided by 3M Health Care and its representatives is intended to provide general information relevant to coding for 3M products. Insurers' reimbursement policies can vary, however, and the use of the codes discussed here does not guarantee that an insurer will cover or pay at any particular level.

References

1. *In-vitro* testing shows that the transparent film dressing provides a viral barrier from viruses 27nm in diameter or larger while the dressing remains intact without leakage.
2. Etris-Brown M, Milne C, Orsted H, Gates J, Netsch D, et al. A prospective, randomized, multisite clinical evaluation of a transparent, absorbent, clear acrylic dressing and a hydrocolloid dressing in the management of stage II and shallow stage III pressure ulcers. *Advances in Skin and Wound Care* 2008;21(4):169-174.
3. LeBlanc K, Baranoski S. Prevention and management of skin tears. *Advances in Skin and Wound Care* 2009;22(7):325-332.
4. Malone ML, Rozario N, Gavinski M, Goodwin J. The epidemiology of skin tears in the institutionalized elderly. *J Am Geriatr Soc* 1991;39(6):591-5.
5. Bank D, Nix D. Preventing skin tears in a nursing and rehabilitation center: an interdisciplinary effort. *Ostomy Wound Management* 2006;52(9):38-46.
6. Gray D, Stringfellow S, Cooper P, Russell F, Johnson S, et al. Pilot RCT of two dressings regimens for the management of skin tears. *Wounds UK* 2011;7(2).
7. Fleck CA. Preventing and treating skin tears. *Advances in Skin and Wound Care* 2007;20(6):315-320.
8. Ratliff CR, Fletcher KR. Skin tears: a review of the evidence to support prevention and treatment. *Ostomy Wound Management* 2007;53(3):32-42.
9. Roberts MJ. Preventing and managing skin tears. *JWOCN* 2007;34(3):256-259.
10. Expert opinion.
11. Cost of RN from Bureau of Labor Statistics (Alternate Care-Average of Long Term Care and Home Care).
12. Skin Tears: State of the Science: Consensus statements for the prevention, predication, assessment and treatment of skin tears. *Advances in Skin & Wound Care* 2011; 24(9) (Suppl 1): 2-15.

3M™ Tegaderm™ Dressings SIMPLE. DEPENDABLE. TRUSTED.

To learn more about 3M™ Tegaderm™ Absorbent Clear Acrylic Dressing, visit us at <http://3M.com/TegadermAbsBrochure>. For more information about the Skin & Wound Care family of products, visit us at www.3M.com/SkinWoundCare, contact your 3M Skin & Wound Care representative or call the 3M Health Care Customer Helpline at 1-800-228-3957. Outside of the United States, contact the local 3M subsidiary.



Skin & Wound Care Division
3M Health Care
 2510 Conway Ave.
 St. Paul, MN 55144
 USA
 1-800-228-3957
www.3M.com/healthcare

3M Canada
 P.O. Box 5757
 London, Ontario N6A 4T1
 Canada
 1-800-364-3577
www.3M.com/ca/healthcare

Mepilex is a registered trademark of Mölnlycke Health Care. Flexigel is a registered trademark of Smith & Nephew. ADAPTIC® is a registered trademark of Systagenix. Telfa is a trademark of Covidien Company.

3M and Tegaderm are trademarks of 3M. Please recycle. Printed in U.S.A. Used under license in Canada. © 3M 2006, 2012. All rights reserved. 70-2009-6371-1